

# NILOTINIB

Most patients treated with Nilotinib will experience adverse effects, but the effects will differ from one patient to the next. Symptoms may indicate that the underlying cancer is not under control or has relapses. Cancer patients may also have co-morbid diseases that require treatment and cause symptoms.

The most common adverse effects with Nilotinib are myelosuppression, rash, pruritus, headache, nausea and vomiting, fatigue, myalgia, nasopharyngitis, constipation, diarrhea, arthralgia, pyrexia (fever), back pain, cough, and asthenia.

### ADVERSE DRUG REACTION MANAGEMENT GUIDE

1. **Myelosuppression**
2. **Nausea & vomiting**
3. **Diarrhea**
4. **Constipation**
5. **Fatigue & weakness**
6. **General pain (headache)**
7. **Joint, muscle, or back pain**
8. **Rash**
9. **Pruritis**

#### 1. Myelosuppression

Nilotinib may cause suppression of the blood cell production in the myeloid tissues of the bone marrow. This can result in lowering of white blood cells and platelets. It is important to have a Complete Blood Count (CBC) blood test prior to the start of each cycle of this agent. If any blood cell component is reduced below an acceptable level, the drug may need to be held until the blood cells recover. **Nilotinib must NOT be dispensed** until the CBC test is completed and verified prior to each cycle of the treatment. Verification will be done by an oncology health professional.

**Prevention:** General infection preventative measures should be followed while on this drug, especially if the blood counts are low. Advise patient to:

- Limit contact with people who are sick, have colds, or have been recently vaccinated
- Rest often
- Do not eat uncooked vegetables
- Wash hands often

If the platelet count is low, tell the patient to take. Advise patient to:

- Take care when shaving or performing any activity of daily living where the skin could be cut
- Use a soft toothbrush.
- Tell your doctor before dental work is done.

**Management:** If the patient has a fever or other signs of an infection when the blood counts are low, advise him/her to go directly to the Emergency Department and contact the oncologist when there. The ER staff needs to be told that the patient is taking this drug, and that it is a form of chemotherapy. Empiric antibiotics will be required.

If the patient has unusual bleeding when the platelet counts are low, advise him/her to go to the Emergency Department, tell the ER staff about this drug, and contact the oncologist when there.

## 2. Nausea & vomiting

Nausea and vomiting may occur in up to 30% of patients on Nilotinib. Unlike the nausea and vomiting often experienced by patients on cytotoxic chemotherapy (acute onset, more emesis than nausea), patients on Nilotinib tend to have nausea of lesser severity and longer duration, with or without emesis. This can be more distressing to patients' quality of life than acute nausea and vomiting. Often patients will have nausea without the relief that comes from emesis.

**Management:** The following may provide relief from nausea and vomiting:

- Prophylactic antiemetic agents (e.g. dopaminergic agents such as prochlorperazine, or promotility agents such as metoclopramide) given with each dose of Nilotinib and repeated as needed for nausea control. While there is no evidence to support the use of dimenhydrinate, there is evidence that ginger products (e.g. Graval<sup>®</sup> Ginger) may be effective, with fewer adverse effects
- Avoid spicy or greasy foods that may contribute to the feeling of nausea. Bland foods, fresh air, and plenty of clear water may reduce the feelings of nausea

## 3. Diarrhea

Diarrhea is a common side effect with Nilotinib, occurring in up to 22% of patients. Dietary modifications are not recommended in anticipation of diarrhea, but must be considered if diarrhea occurs.

**Management:**

For mild diarrhea (less than 4 loose stools per day)

- Follow instructions on loperamide (e.g., Imodium<sup>®</sup>) package insert: 2 tablets immediately, then 1 tablet after each liquid bowel movement (maximum: 8 tablets/24 hours)

For moderate diarrhea (more than 4 to 6 loose stools per day or night-time diarrhea), tell the patient to be more aggressive with loperamide (e.g., Imodium<sup>®</sup>) for early-onset diarrhea

- Take 2 tablets immediately, then 1 tablet every 2 hours during the day and 2 tablets every 4 hours during the night until bowel movements are normal for at least 12 hours
- This dosage is higher than packaging recommendations.

Replace lost fluids: Fluid intake is more important than eating in patients with diarrhea. To replace lost fluid, advise patients to increase fluids by up to 3 to 4 liters per day (unless there is a known contraindication to increased fluid intake). The patient may drink several types of fluid, including plain water and electrolyte-containing drinks, such as clear broth, gelatin desserts, sports drinks, flat soft drinks, or decaffeinated tea

Anal care: Recommend to your patient to:

- Clean the anal area with mild soap and warm water after each bowel movement to prevent irritation
- Apply a barrier cream or ointment, such as petroleum jelly or Isle's paste
- Soak in a warm bathtub or sitz bath to relieve discomfort

Dietary changes during diarrhea: Advise your patients to change their diet while diarrhea is a problem:

- Eat and drink small quantities of food often
- Avoid spicy, greasy, or fried foods
- Follow the BRAT (banana, rice, applesauce, toast) diet, along with clear liquids, until diarrhea begins to resolve
- Follow a lactose-free diet
- Avoid cabbage, brussel spouts, and broccoli, which may produce stomach gas, bloating and cramps

#### **4. Constipation**

Constipation is generally understood to be a reduction in bowel movements to fewer than 3 per week, or unsatisfactory (e.g. incomplete) defecation. Reduced bowel function may give the misperception of constipation to patients who expect one or more movements daily. Constipation may be caused by this medication, other medications being taken concurrently (e.g. narcotic analgesics), or by the underlying cancer. Reduced physical activity or dietary changes, possibly related to this medication or the overall cancer treatment, may contribute to constipation. Abdominal pain is often associated with constipation, but if symptoms become severe a medical assessment to examine for fecal impaction or bowel obstruction may be needed.

**Prevention:** Advise patient to:

- Change diet, if possible, to include more fruits, vegetables, and high-fibre foods. If the patient is taking a low-calorie diet, consider increasing calories to improve colonic transit.
- Use of a regular laxative regimen, such as senna or bisacodyl at bedtime
- Consider a bowel routine to maintain regularity, such as attempting bowel movements each morning after breakfast (usually the optimal time for a movement), using the toilet when there is an urge instead of repressing it, placing a footstool in front of the toilet to elevate thighs during movement.
- Consider adding light exercise for patients with mostly sedentary lifestyles
- Weight loss in over-weight patients may help improve bowel function

**Management:**

- If prophyllactive stimulant laxative is not effective, try osmotic laxative (e.g. lactulose or PEG)
- Stool softeners have not been shown to be effective
- Drink of fluids

Biofeedback and relaxation techniques may help some patients with pelvic floor dysfunction.

Psychosocial teams in the cancer centres may help with this type of intervention.

## 5. Fatigue & weakness

About 15-30% of patients on Nilotinib will experience fatigue. Fatigue is not life-threatening but will significantly reduce quality of life.

**Management:** The following may provide relief from fatigue:

- There are no medications that have demonstrated an effect to relieve fatigue
- Mild exercise is very helpful to reduce fatigue, but must be manageable if there is also muscle weakness

## 6. General pain (headache)

Patients on Nilotinib may experience other types of pain. About 20-30% of patients have headaches while on this treatment. Generalized pain may be a drug side effect or may be related to the cancer.

**Management:** The following may provide relief from headaches and other general pain problems:

- Mild pain may respond to non-pharmacologic approaches, such as rest, distraction, cool cloth on the forehead
- Mild pain medications- Acetaminophen preferred; do NOT use non-steroidal anti-inflammatory agents, prescription or OTC due to risk of bleeding (e.g. ibuprofen, ASA, naproxen)
- Acetaminophen with codeine, either OTC (low dose) or on prescription (higher dose) may be considered for more severe pain.
- If acetaminophen is not sufficient to control pain, consider prescription opioid analgesics for management of more severe pain (possibly due to tumor)
- If there is a neuropathic component to the pain, consider a trial with a tricyclic antidepressant (e.g. low dose amitriptyline or imipramine) or gabapentin

## 7. Joint, muscle, or back pain

Aching backs or muscles and muscle cramps occur in 6-8% of patients taking Nilotinib. Muscle cramps may occur in the hands, feet, calves, or thighs. Cramps have been described as sustained muscular contractions that follow a consistent pattern, frequency, and severity. Muscle cramps may be related to exertion or could happen at night. Patients should avoid using quinine or drinking tonic water (contains quinine).

Joint pain may begin in the first month of therapy and commonly subside after a few months. Pain may affect the leg bones, hips, and knees, and may appear in an asymmetrical pattern. Although there are no evidence-based guidelines for prevention or treatment, anecdotal reports and expert experience suggest that some patients' pain could be eased by using mineral supplements.

**Management:** The following may provide relief from muscle aches or cramps:

- Calcium and magnesium supplements
- Mild pain medications- Acetaminophen preferred; do NOT use non-steroidal anti-inflammatory agents, prescription or OTC due to risk of bleeding (e.g. ibuprofen, ASA, naproxen)

- For bone or joint pain, consider the use of heating pads, ice packs, or topical arthritis creams and liniments
- Mild exercise and/or massage therapy may help reduce bone and joint pain
- Avoid using quinine or drinking tonic water

## 8. Rash

Rash is a common adverse effect of Nilotinib, occurring 20% of patients. Rash symptoms often appear soon after starting treatment. This rash presents with spots and bumps on the forearms, trunk, and sometimes, the face. They are often itchy, but if scratched, may become infected and crusty. Most cases of this generalized skin rash are mild and go away on their own. Rash is more common in women and patients on higher doses, and may worsen after sun exposure.

It is important to recognize rash symptoms early and start symptomatic therapy promptly.

**Prevention:** Prevention should begin when Nilotinib therapy is begun, and continue throughout treatment.

You should advise your patient to:

- Cleanse with mild soaps or hypoallergenic cleaners or bath or shower oils to avoid skin dryness
- Take short showers with warm water.
- Moisturize twice a day with a colloidal oatmeal lotion, such as Aveeno® lotion, or thick, emollient-based creams, such as Neutrogena® Norwegian Formula hand cream or Vaseline Intensive Care® Advanced Healing Lotion
- Use only fragrance-, alcohol-, and dye-free lotions and cosmetics
- Use a dermatologist-approved cover-up, such as Dermablend® or Cover FX®
- Remove make-up with a gentle, skin-friendly cleanser (e.g., Neutrogena®, Dove®).
- Use a broad-spectrum sunscreen (SPF of 30 or more) that contains zinc oxide or titanium dioxide

**Management:** For Mild to moderate skin rash, there are some over-the-counter options you may consider:

- Antihistamine (diphenhydramine)
- Topical steroid (hydrocortisone 0.5%)
- Coal tar preparations

If the rash progresses to moderate to severe, the patient may need prescribed therapy:

- Oral corticosteroids (short course, with or without topical triamcinolone acetonide 0.1% ointment)
- Temporary interruption of therapy until the rash resolves, and then re-challenge at low dose

### Counseling tips:

Reinforcement is important. Make a note to yourself (or book a time for a follow up call) to repeat these suggestions 2-3 weeks after the Nilotinib treatment initiation or any report of early rash symptoms.

## 9. Pruritis

Pruritus (itchiness) happens in some patients, and usually occurs because skin has lost its moisture.

**Prevention:** Preventing dry skin is the key to preventing pruritus. Advise your patients to:

- Use mild soaps that are deodorant and fragrance-free (e.g. Dove® or Neutrogena®)
- Apply lotions or bland emollients (Eucerin® cream, Neutrogena® Norwegian Formula Hand Cream, Vaseline Intensive Care® Advanced Healing Lotion) often.
- Choose “anti-itch” products
- Use liquid shower gels in place of soap.

**Management:** For mild to moderate pruritus, consider advising patients to:

- Apply more lotion than usual to help eliminate itchiness
- Use lotions that contain aloe vera or dimethicone Moisturel®
- Use antidandruff shampoo and conditioner
- Use hair products that contain tea tree oil, which contain extra moisturizers and may help with symptoms

For moderate to severe pruritus with intense, widespread itching, refer patient to physician.

Antihistamines may provide some relief.

#### REFERENCES:

ONTarget Resource Guide, Common Side Effects from Targeted Therapy. The Groupe d'étude en oncologie and The Canadian Association of Pharmacy in Oncology, 2012.

Systemic Therapy Manual for Cancer Treatment, Cancer Care Nova Scotia, 2013

Tasigna® Product Monograph, Novartis Pharmaceuticals Corp., June 2010

Patient Self-Care. Helping Your Patients Make Therapeutic Choices. Canadian Pharmaceutical Association, 2010

Therapeutic Choices, Sixth Edition, Canadian Pharmaceutical Association, 2011