

Guidelines for ordering Chest X-Rays - PA and Lateral or Portable

Not indicated	Indicated
Asymptomatic patients:	Symptoms or signs of a new cardiothoracic illness (Eg. Acute cardiac respiratory symptoms in a patient
	with no recent or accessible chest x-ray)
Upper respiratory tract infection	Lower respiratory tract infection involves:BronchiLungs
Pneumonia patients who are improving but aren't: COPD patients High cancer risk Chronic smokers >55 years old	Pneumonia patients if new symptoms occur, or if patient: • Has COPD • Is a chronic smoker • Has high cancer risk • Is >55 years old Repeat Chest X-Ray should be no earlier than 6 weeks after last X-Ray
Routine chest X-Ray's in: ICU Post-Op PACU Known chronic lung disease with acute shortness of	Each Chest X-Ray requires a clinical indication Need a clinical concern for a co-existent complication
breath	Need a clinical concern for a co-existent complication
Known coronary disease with symptoms of patient's usual chest pain	Atypical new painConcern for complication
 Minor trauma Injury severity score < 15 Stable vital signs No internal organ damage No major bleeding 	Major trauma Fall > 20 ft. Car - pedestrian or motorcycle accident MVA > 40 mph Ejection or Prolonged extraction from vehicle Other death in vehicle Chronic steroid use
Lung cancer screening	Lung cancer screening requires a CT and only if patient is: High risk 30 pkg / years +ve Family Hx Had a previous cancer <75 years old
Thoracic aneurysm F/U	Aneurysms require a CT

June 2013